

Rapid Communication

## The neuroanatomy of general intelligence: sex matters

Richard J. Haier,<sup>a,\*</sup> Rex E. Jung,<sup>b</sup> Ronald A. Yeo,<sup>c</sup> Kevin Head,<sup>a</sup> and Michael T. Alkire<sup>d</sup>

<sup>a</sup>Department of Pediatrics, University of California, Med. Sci. I, B140, Irvine, CA 92697-5000, USA

<sup>b</sup>Department of Neurology and MIND Institute, University of New Mexico, Albuquerque, NM 87131, USA

<sup>c</sup>Department of Psychology, University of New Mexico, Albuquerque, NM 87131, USA

<sup>d</sup>Department of Anesthesiology, University of California Irvine Medical Center, Bldg. 53, 104B, Orange, CA 92868-1350, USA

Received 21 October 2004; revised 4 November 2004; accepted 9 November 2004

Available online 16 January 2005

**We examined the relationship between structural brain variation and general intelligence using voxel-based morphometric analysis of MRI data in men and women with equivalent IQ scores. Compared to men, women show more white matter and fewer gray matter areas related to intelligence. In men IQ/gray matter correlations are strongest in frontal and parietal lobes (BA 8, 9, 39, 40), whereas the strongest correlations in women are in the frontal lobe (BA10) along with Broca's area. Men and women apparently achieve similar IQ results with different brain regions, suggesting that there is no singular underlying neuroanatomical structure to general intelligence and that different types of brain designs may manifest equivalent intellectual performance.**

© 2004 Elsevier Inc. All rights reserved.

*Keywords:* Neuroanatomy; General intelligence; Sex

### Introduction

Recent evidence from structural brain imaging of normal adults indicates that individual differences in general intelligence as assessed with the Wechsler Adult Intelligence Scale (WAIS), one measure of the intelligence quotient (IQ), are strongly related to differences in gray matter (GM) and white matter (WM) volumes in a number of specific areas distributed mostly in frontal, temporal, and parietal regions (Haier et al., 2004). Moreover, the relative contribution of GM and WM volumes in these areas to explaining variance in intelligence scores shifts with age. For instance, older adults show more GM–IQ relationships in the frontal and parietal areas, whereas younger adults show more relationships in temporal and limbic areas.

Comparisons of general intelligence assessed with standard measures like the WAIS show essentially no differences between men and women (Jensen, 1998), but it remains to be determined

whether the neuroanatomical basis for general intelligence is the same in men and women. A myriad of sex differences in brain structure have been reported (de Courten-Myers, 1999; Luders et al., 2004). Here, we examine whether there are male and female differences in the correlations between IQ and GM or WM volumes based on optimized voxel-based morphometry (VBM) of structural MRI data. Earlier studies of brain structure and intelligence were based on region-of-interest (ROI) methods, which have a number of limitations for examining the entire brain (Andreasen et al., 1993; Flashman et al., 1997). Functional brain imaging studies have reported a number of sex differences (Cahill et al., 2001; Gur et al., 2000; Haier and Benbow, 1995; Mansour et al., 1996; Neubauer et al., 2002; Shaywitz et al., 1995, 2001), but task-specific demands on cognitive resources with functional studies must always be considered as the task itself could affect interpretation of functional imaging results. Structural imaging correlated with off-line analyses of various cognitive performance measures and traits, on the other hand, can identify those differences in neuroanatomy which may underlie the cognitive measure of interest, irrespective of any task design constraints.

### Materials and methods

#### Subjects

Two independent samples of normal volunteers were studied. The first sample of 23 volunteers (14 women and 9 men; mean age = 27, SD = 5.9, range = 18–37) were recruited from the University of New Mexico (UNM). The second sample of 25 volunteers (13 men and 12 women, mean age = 59, SD = 16; range 37–84) were recruited from the University of California, Irvine (UCI) as middle-aged and older normal controls for an imaging study of dementia in Down syndrome and Alzheimer's disease (Haier et al., 2003a). These are the same subjects used in our previous report (Haier et al., 2004), except that one additional female subject was added to the UCI sample. All of the 48 subjects were in good physical and mental health; none had a history of head injury and none showed any clinical signs of dementia.

\* Corresponding author. Fax: +1 949 854 1989.

E-mail address: rjhaier@uci.edu (R.J. Haier).

Available online on ScienceDirect (www.sciencedirect.com).

### Intelligence testing

To assess general intelligence, subjects were tested with the WAIS. The WAIS battery (Wechsler, 1981) consists of 11 diverse subtests which tap a variety of verbal and non-verbal mental abilities which contribute to general intelligence. The WAIS full scale IQ score (FSIQ) is based on performance of all 11 subtests (according to age based norms). Factor analytic studies (Jensen, 1998) show that each subtest loads on the *g* factor and the FSIQ score loads the highest (about 0.90 or 81% of the variance in *g*). For this reason, FSIQ is considered one of the best indexes of individual differences in general intelligence, as first described by Spearman (1904). Mean FSIQ for all 23 males was 118.8 (SD = 15.1; range 95–155) and 113.5 (SD = 13.4; range 90–134) for all 25 females. This was not significantly different between groups ( $t = 1.29$ ,  $P = 0.20$ ; two-tailed). In sample 1 (UNM), mean FSIQ for the males was 122.8 (SD = 16; range 109–155) and 111.1 (SD = 12.4; range 90–133) for females. In sample 2 (UCI), mean FSIQ for the males was 116.3 (SD = 14.5; range 95–142) and 116.5 (SD = 14.6; range 90–134) for females.

Six of the WAIS subtests can be summarized as a single score on verbal IQ (VIQ) and five subtests can be summarized as performance IQ (PIQ). These are highly inter-correlated ( $r = 0.83$  in the male sample;  $r = 0.68$  in the female sample) and with FSIQ (0.97 and 0.94, respectively, in the males; 0.96 and 0.86 in the females). Although FSIQ is the single best summary score for general intelligence, VIQ and PIQ scores may yield additional information when correlated to VBM results.

### Structural MRI imaging parameters

Sample 1 MRIs were obtained with a 1.5-T scanner, head coil, and software (Signa 5.4; General Electric Medical Systems, Waukesha, WI). A T1 sagittal localizer sequence (TE = 6.9 ms, TR = 200 ms, FOV =  $24 \times 24$  cm<sup>2</sup>, five slices, thickness = 5 mm, spacing = 2.5 mm, matrix =  $256 \times 128$ ) was acquired, followed by a T1 weighted axial series (fast RF spoiled gradient-recalled, TE = 6.9 ms, TR = 17.7 ms, flip angle = 25°, matrix =  $256 \times 192$ , 120 slices, thickness = 1.5 mm) to give full brain coverage. Sample 2 MRIs were obtained with a 1.5-T clinical Phillips Eclipse scanner (Philips Medical Systems, N.A., Bothell, WA). We used T1-weighted, volumetric SPGR MRI scans (FOV = 24 cm, flip angle = 40, TR = 24, TE = 5). The images consisted of 120 contiguous 1.2-mm-thick axial slices, each with an in-plane image matrix of  $256 \times 256$  image elements, to achieve full brain coverage. All images in both samples were visually inspected to ensure image quality.

### Voxel-based morphometry (VBM)

We applied VBM to identify brain areas where GM and WM volumes were correlated to IQ separately for males and females. We used Statistical Parametric Mapping software (SPM2; The Wellcome Department of Imaging Neuroscience, University College London) to create one study-specific template for the combined sample of UNM and UCI males and a separate template for the combined female samples. The optimized VBM protocol was applied to males and females separately using the methods of Ashburner and Friston (2000) and Good et al. (2001). To preserve the amount of tissue in any given anatomical region after spatial normalization, the optimal GM and WM partitions were multiplied by the Jacobian determinants of their respective spatial trans-

formation matrix. The reason for performing this modulation step is so that the final VBM statistics will reflect local deviations in the absolute amount (volume) of tissue in different regions of the brain (Ashburner and Friston, 2000). Since this was done separately for the men and women, brain size differences between the groups should not be confounded. The modulated GM and WM partitions were then smoothed with a 12-mm FWHM isotropic Gaussian kernel to account for slight misalignments of homologous anatomical structures and to ensure statistical validity under parametric assumptions.

### Statistical conjunction approach

We specifically tested whether regional GM or WM volumes were correlated with FSIQ scores treating any effects of age and intra-sex brain size (total brain GM or WM, respectively) as nuisance variables in the SPM2 design matrix. We used the conjunction approach (Price and Friston, 1997) to show where GM and WM correlations with FSIQ overlapped for the UNM and the UCI samples (i.e., voxels with correlations in common for both samples) separately for the males and females. We repeated these analyses separately for VIQ and PIQ. The conjunction approach minimizes potential problems associated with combining data from different scanners and has the additional advantage of maximizing statistical power because all subjects are used in the analysis. Findings are considered significant at  $P < 0.05$  corrected for multiple comparisons; findings at  $P < 0.001$  uncorrected also are shown for hypothesis generation regarding FSIQ.  $R^2$  estimates were determined using the formula:  $R^2 = t^2 / (df + t^2)$ , where  $df = 17$  for the males and 19 for the females (from the SPM2 conjunction design matrix). Locations of significant clusters (centroids) are converted from Montreal Neurological Institute (MNI) to Talairach atlas (Talairach and Tournoux, 1988) coordinates and reported as closest Brodmann area (BA) where possible. Only clusters of at least 10 voxels are reported.

### Results

In our sample of men, the conjunction analysis showed GM volume was correlated to FSIQ most strongly ( $P < 0.05$  corrected for multiple comparisons) in bilateral frontal lobes (BA 8, 9) and in left parietal lobe; Wernicke's area (BA 39 and 40). In the women, the strongest GM-FSIQ correlation was in the right frontal lobe (BA10) and the largest cluster was in Broca's area (left BA 44, 45). These significant areas are shown in Fig. 1; the exact anatomical locations and cluster sizes are listed in Table 1, along with the estimated variance accounted for in FSIQ by each GM cluster (41–54% for the most significant clusters).

WM volume correlations with FSIQ were not as strong ( $P < 0.001$ , uncorrected), but they were more extensive in the women, as illustrated in Fig. 1 and detailed in Table 2. All correlations in Tables 1 and 2 are positive; there were no appreciable negative correlations. As shown in Table 3, most of the GM findings for men were evenly distributed in frontal and parietal lobes (45% each), whereas the GM findings were predominately in the frontal lobe (84%) for women. WM findings were predominately in temporal lobes for the men (82%) and in frontal lobes for the women (86%).

GM voxels correlated to FSIQ represented 9.6% of all GM voxels (693,360) in the male brain. GM voxels correlated to FSIQ




Fig. 1. Correlations ( $P < 0.001$ ) in men ( $N = 23$ ) and women ( $N = 25$ ) between gray matter and white matter volumes and IQ conjuncted across the UNM and the UCI samples. Anatomical locations, atlas coordinates and cluster sizes are listed in Tables 1 and 2. Color scale bars at bottom refer to  $t$  test values for each analysis.

represented only 1.7% of all GM voxels (612,007) in the female brain. WM voxels correlated to FSIQ represented only 0.1% of all WM voxels (492,814) in the male brain. WM voxels correlated to FSIQ represented 1.3% of all WM voxels (416,630) in the female brain.

VIQ correlations with GM and WM were computed using PIQ as an additional nuisance variable along with age and total GM (or WM); the PIQ correlations used VIQ as an additional nuisance variable. Only two results were significant when corrected for multiple comparisons. In the men, PIQ correlated with GM ( $P = 0.057$ ) in BA7 (right parietal lobe;  $x = 10, y = -57, z = 59$ ). In the women, VIQ correlated with GM ( $P = 0.018$ ) in the thalamus/pulvinar area ( $x = 4, y = -29, z = 15$ ).

## Discussion

These results expand upon our previous report (Haier et al., 2004), which combined men and women into a single analysis to maximize statistical power. In that analysis, GM and WM volumes were correlated to FSIQ with sex treated as a nuisance variable; the

results showed correlations in several regions distributed throughout the brain. Here, we have systematically sought to explore sex differences in the structural brain organization underlying general intelligence. Of the major areas identified in our earlier analysis combining men and women, the sex-specific analyses show that some of the previous significant areas are found more in men and others are found more in the women; the relative roles of GM and WM also differ between the men and women.

These present results highlight an important dissociation of brain morphology related to intellectual functioning in normal adult brains, as the pattern of voxel types and voxel locations linked with intellectual functioning differed substantially between the sexes. With respect to voxel types, men had roughly 6.5 times the number of GM voxels identified as related to intellectual functioning as did women, and women had roughly nine times more WM voxels than did men. With respect to regional effects, in women, 84% of the identified GM voxels correlated to IQ were in the frontal region, as compared to 45% in men. Even greater sex differences were observed in WM, where for women 86% of the identified voxels were frontal, as compared to 0% in men. More left hemisphere voxels (GM plus WM) were identified in both men

Table 1

Areas of Correlation ( $P < 0.001$ ) between gray matter and IQ in males (top) and females (bottom) conjuncted for UNM and UCI samples (see Fig. 1)

| Nearest gray matter                          | Cluster size | x   | y   | z   | t    | $r^2$ |
|--|--------------|-----|-----|-----|------|-------|
| <i>Males</i>                                 |              |     |     |     |      |       |
| Right superior frontal gyrus, BA 8*          | 28,757       | 31  | 43  | 40  | 4.51 | 0.54  |
| Right superior frontal gyrus, BA 9*          |              | 8   | 55  | 26  | 4.29 | 0.52  |
| Left superior frontal gyrus, BA 9*           |              | -5  | 53  | 28  | 4.09 | 0.50  |
| Left parietal lobe, angular gyrus, BA 39*    | 16,938       | -51 | -59 | 34  | 4.09 | 0.50  |
| Left temporal, supramarginal gyrus, BA 40*   |              | -61 | -50 | 25  | 3.84 | 0.46  |
| Left superior temporal gyrus, BA 39*         |              | -56 | -57 | 28  | 3.81 | 0.46  |
| Right parietal lobe, precuneus, BA 39        | 11,394       | 44  | -72 | 36  | 3.37 | 0.40  |
| Right inferior parietal lobule, BA39         |              | 45  | -64 | 41  | 3.23 | 0.38  |
| Right superior temporal gyrus, BA 42         |              | 69  | -30 | 21  | 3.20 | 0.38  |
| Left inferior temporal gyrus, BA 20          | 886          | -55 | -8  | -32 | 3.33 | 0.39  |
| Left inferior temporal gyrus, BA 20          |              | -49 | -1  | -32 | 2.30 | 0.24  |
| Right parietal lobe, precuneus, BA 31        | 1724         | 19  | -67 | 26  | 3.33 | 0.39  |
| Right parietal lobe, precuneus, BA 18        |              | 18  | -68 | 18  | 2.19 | 0.22  |
| Right middle occipital gyrus, BA 18          | 3030         | 43  | -79 | -10 | 3.16 | 0.37  |
| Right inferior occipital gyrus, BA 18        |              | 35  | -87 | -6  | 3.01 | 0.35  |
| Right middle occipital gyrus, BA 19          |              | 27  | -84 | 20  | 2.45 | 0.26  |
| Left superior frontal gyrus, BA 8            | 439          | -41 | 21  | 51  | 2.65 | 0.29  |
| Left middle frontal gyrus, BA 8              |              | -40 | 27  | 42  | 2.28 | 0.23  |
| Left middle frontal gyrus, BA 8              |              | -34 | 31  | 45  | 2.11 | 0.21  |
| Left inferior temporal gyrus, BA 37          | 472          | -48 | -71 | -2  | 2.48 | 0.27  |
| Left middle occipital gyrus, BA 18           | 1050         | -21 | -97 | 7   | 2.43 | 0.26  |
| Left middle occipital gyrus, BA 18           |              | -23 | -90 | 19  | 2.37 | 0.25  |
| Left middle occipital gyrus, BA 18           |              | -28 | -93 | 8   | 2.24 | 0.23  |
| Right occipital lobe, lingual gyrus, BA 18   | 552          | 10  | -74 | -2  | 2.42 | 0.26  |
| Right occipital lobe, lingual gyrus, BA 18   |              | 6   | -78 | -7  | 2.16 | 0.22  |
| Left middle frontal gyrus, BA 9              | 116          | -50 | 16  | 36  | 2.40 | 0.25  |
| Left superior frontal gyrus, BA 9            | 211          | -24 | 51  | 32  | 2.38 | 0.25  |
| Left superior frontal gyrus, BA 8            |              | -25 | 44  | 39  | 2.24 | 0.23  |
| Left sub-lobar extra-nuclear, claustrum      | 61           | -23 | 28  | 4   | 2.38 | 0.25  |
| Left sub-lobar extra-nuclear, claustrum      |              | -22 | 22  | 11  | 2.22 | 0.22  |
| Left superior frontal gyrus, BA 8            | 450          | -21 | 26  | 52  | 2.34 | 0.24  |
| Left superior frontal gyrus, BA 8            |              | -16 | 33  | 52  | 2.11 | 0.21  |
| Left middle frontal gyrus, BA 6              |              | -26 | 20  | 51  | 2.09 | 0.20  |
| Right medial frontal gyrus, BA 6             | 151          | 1   | -9  | 62  | 2.27 | 0.23  |
| Left inferior occipital gyrus, BA 18         | 42           | -34 | -85 | -17 | 2.23 | 0.23  |
| Right sub-lobar thalamus                     | 24           | 22  | -22 | 18  | 2.19 | 0.22  |
| Left middle frontal gyrus, BA 10             | 12           | -39 | 54  | 10  | 2.18 | 0.22  |
| Left superior temporal gyrus, BA 22          | 15           | -57 | -17 | 1   | 2.16 | 0.22  |
| Left temporal lobe, fusiform gyrus, BA19     | 28           | -46 | -78 | -14 | 2.14 | 0.21  |
| Left sub-lobar, thalamus, pulvinar           | 29           | -26 | -28 | 12  | 2.12 | 0.21  |
| Right occipital lobe, cuneus, BA 19          | 16           | 6   | -78 | 37  | 2.09 | 0.20  |
| Right inferior frontal gyrus, BA 47          | 41           | 13  | 10  | -23 | 2.09 | 0.20  |
| Left occipital lobe, lingual gyrus, BA 18    | 18           | -11 | -76 | 2   | 2.06 | 0.20  |
| <i>Females</i>                               |              |     |     |     |      |       |
| Right middle frontal gyrus, BA 10*           | 2025         | 40  | 40  | 13  | 3.61 | 0.41  |
| Right middle frontal gyrus, BA 10            |              | 35  | 37  | 7   | 3.21 | 0.35  |
| Left inferior frontal gyrus, BA 45           | 4663         | -49 | 36  | 7   | 2.78 | 0.29  |
| Left inferior frontal gyrus, BA 46           |              | -46 | 41  | 2   | 2.66 | 0.27  |
| Left inferior frontal gyrus, BA 47           |              | -52 | 28  | 0   | 2.61 | 0.26  |
| Left limbic lobe, cingulate gyrus, BA 31     | 446          | -15 | -40 | 30  | 2.66 | 0.27  |
| Right middle frontal gyrus, BA 9             | 907          | 40  | 18  | 33  | 2.64 | 0.27  |
| Right middle frontal gyrus, BA 9             |              | 35  | 29  | 32  | 2.30 | 0.22  |
| Left parietal lobe, postcentral gyrus, BA 40 | 141          | -67 | -18 | 15  | 2.61 | 0.26  |
| Left inferior frontal gyrus, BA 11           | 632          | -29 | 28  | -22 | 2.59 | 0.26  |
| Left inferior frontal gyrus, BA 11           |              | -34 | 36  | -9  | 2.36 | 0.23  |
| Left middle frontal gyrus, BA 11             |              | -31 | 36  | -17 | 2.33 | 0.22  |
| Left superior frontal gyrus, BA 10           | 103          | -12 | 65  | 23  | 2.59 | 0.26  |
| Left limbic lobe, anterior cingulate, BA 32  | 344          | -15 | 27  | 24  | 2.51 | 0.25  |
| Left middle frontal gyrus, BA 8              | 109          | -32 | 35  | 45  | 2.43 | 0.24  |
| Right middle temporal gyrus, BA 21           | 404          | 56  | -40 | -5  | 2.38 | 0.23  |
| Right middle temporal gyrus, BA 22           |              | 57  | -40 | 2   | 2.19 | 0.20  |

(continued on next page)

Table 1 (continued)

| Nearest gray matter                        | Cluster size | <i>x</i> | <i>y</i> | <i>z</i> | <i>t</i> | <i>r</i> <sup>2</sup> |
|--|--------------|----------|----------|----------|----------|-----------------------|
| <i>Females</i>                             |              |          |          |          |          |                       |
| Left inferior occipital gyrus, BA 19       | 143          | −32      | −77      | −4       | 2.38     | 0.23                  |
| Right superior frontal gyrus, BA 8         | 45           | 39       | 26       | 51       | 2.36     | 0.23                  |
| Right medial frontal gyrus, BA 10          | 104          | 14       | 51       | 4        | 2.29     | 0.22                  |
| Left inferior temporal gyrus, BA 20        | 45           | −67      | −11      | −23      | 2.24     | 0.21                  |
| Left medial frontal gyrus, BA 6            | 151          | 0        | 39       | 35       | 2.21     | 0.20                  |
| Right superior frontal gyrus, BA 6         | 20           | 11       | 31       | 58       | 2.19     | 0.20                  |
| Left middle temporal gyrus, BA 21          | 132          | −63      | −45      | −4       | 2.18     | 0.20                  |
| Right inferior frontal gyrus, BA 47        | 53           | 55       | 38       | −7       | 2.16     | 0.20                  |
| Left frontal lobe, precentral gyrus, BA 44 | 73           | −49      | 15       | 7        | 2.16     | 0.20                  |
| Left medial frontal gyrus, BA 8            | 39           | −8       | 33       | 37       | 2.15     | 0.20                  |
| Left medial frontal gyrus, BA 10           | 19           | 0        | 64       | 12       | 2.15     | 0.20                  |
| Left middle frontal gyrus, BA 10           | 48           | −31      | 52       | 0        | 2.13     | 0.19                  |
| Right superior frontal gyrus, BA 6         | 27           | 14       | 19       | 65       | 2.11     | 0.19                  |
| Right superior frontal gyrus, BA 6         |              | 5        | 17       | 63       | 2.11     | 0.19                  |
| Left sub-lobar insula, BA 13               | 10           | −33      | −31      | 20       | 2.02     | 0.18                  |

*x*, *y*, *z* coordinates converted from original MNI to Talairach.

\* *P* < 0.05 corrected.

and women (64.6% and 65.3%, respectively), consistent with previous reports of left lateralization of both anatomic and higher brain functions (Geschwind, 1972).

It is well established that men have a larger cerebrum than women by about 8–10% (Filipek et al., 1994; Nopoulos et al., 2000; Passe et al., 1997a,b; Rabinowicz et al., 1999; Witelson et al., 1995). Thus, for the current study, we created separate templates for males and females to facilitate comparisons between the sexes, as opposed to “stretching” one gender’s brain to fit the size of another. Many studies have now confirmed a consistent sex difference in human brains: the GM/WM ratio is slightly higher in women than men (Allen et

al., 2003; Goldstein et al., 2001; Gur et al., 1999; Passe et al., 1997a,b; Peters et al., 1998; Schlaepfer et al., 1995). In our sample, using the voxel classifications obtained with VBM, women showed a slightly higher GM/WM matter ratio (1.47) than did men (1.41), consistent with these previous reports. More specific regional analyses of GM volume differences (Goldstein et al., 2001) have shown that women have greater GM volume than men in regions including the precentral gyrus (BA 6, 4), fronto-orbital cortex (BA 47), superior frontal (BA 6, 8, 9), and lingual gyri (BA 17, 18), while men had larger volumes in frontomedial cortex (BA 11, 12), hypothalamus, amygdala, and angular gyrus (BA 39).

Table 2

Areas of Correlation (*P* < 0.001) between white matter and IQ in males (top) and females (bottom) conjuncted for UNM and UCI samples (see Fig. 1)

| Nearest gray matter                   | Cluster size | <i>x</i> | <i>y</i> | <i>z</i> | <i>t</i> | <i>r</i> <sup>2</sup> |
|---------------------------------------|--------------|----------|----------|----------|----------|-----------------------|
| <i>Males</i>                          |              |          |          |          |          |                       |
| Left middle temporal gyrus, BA 37     | 498          | −50      | −56      | 4        | 2.69     | 0.30                  |
| Left inferior temporal gyrus, BA 37   |              | −45      | −52      | −1       | 2.59     | 0.28                  |
| Left parietal, sub-gyral WM matter    | 90           | −28      | −51      | 31       | 2.21     | 0.22                  |
| Right middle temporal gyrus, BA 37    | 19           | 49       | −55      | 5        | 2.18     | 0.22                  |
| Left parietal lobe, sub-gyral, BA 37  | 25           | −24      | −59      | 28       | 2.12     | 0.21                  |
| <i>Females</i>                        |              |          |          |          |          |                       |
| Left middle temporal gyrus, BA 20     | 380          | −51      | −37      | −12      | 3.46     | 0.39                  |
| Left medial frontal gyrus, BA 10      | 1487         | −16      | 54       | 2        | 3.36     | 0.37                  |
| Left medial frontal gyrus, BA 10      |              | −8       | 64       | 0        | 2.15     | 0.20                  |
| Left medial frontal gyrus, BA 10      |              | −17      | 47       | −8       | 2.11     | 0.19                  |
| Left inferior frontal gyrus, BA 47    | 1652         | −47      | 37       | −9       | 3.34     | 0.37                  |
| Left inferior frontal gyrus, BA 47    |              | −36      | 31       | 1        | 2.97     | 0.32                  |
| Left middle frontal gyrus, BA 47      |              | −43      | 33       | −4       | 2.89     | 0.31                  |
| Right medial frontal gyrus, BA 9      | 722          | 4        | 45       | 16       | 3.32     | 0.37                  |
| Right medial frontal gyrus, BA 9      |              | 2        | 53       | 19       | 2.88     | 0.30                  |
| Right middle temporal gyrus, BA 21    | 106          | 56       | −13      | −7       | 2.74     | 0.28                  |
| Right superior frontal gyrus, BA 10   | 997          | 27       | 54       | 7        | 2.73     | 0.28                  |
| Right medial frontal gyrus, BA 10     |              | 23       | 47       | 11       | 2.45     | 0.24                  |
| Left superior parietal lobule, BA 7   | 205          | −24      | −50      | 60       | 2.66     | 0.27                  |
| Left transverse temporal gyrus, BA 41 | 15           | −42      | −23      | 14       | 2.33     | 0.22                  |
| Left middle temporal gyrus, BA 22     | 26           | −55      | −40      | 3        | 2.24     | 0.21                  |
| Left sub-lobar, insula, BA 13         | 20           | −44      | −11      | 20       | 2.06     | 0.18                  |
| Left sub-lobar, insula, BA 13         | 11           | −44      | −17      | 21       | 2.05     | 0.18                  |

*x*, *y*, *z* coordinates converted from original MNI to Talairach.

Table 3

Number and size of voxel clusters with a significant correlation ( $P < 0.001$ ) between gray and white matter volume and IQ for each brain lobe in the male ( $N = 23$ ) and female ( $N = 25$ ) samples<sup>a</sup>

|                        | Frontal | Parietal | Temporal | Limbic | Occipital | Sub-lobar | Totals |
|------------------------|---------|----------|----------|--------|-----------|-----------|--------|
| <i>Gray matter</i>     |         |          |          |        |           |           |        |
| Male no. of clusters   | 8       | 3        | 4        | 0      | 6         | 3         | 24     |
| Male total voxels      | 30,177  | 30,056   | 1401     | 0      | 4708      | 114       | 66,456 |
| % of total voxels      | 45      | 45       | 2        | –      | 7         | <1        |        |
| Female no. of clusters | 16      | 1        | 3        | 2      | 1         | 1         | 24     |
| Female total voxels    | 9018    | 141      | 581      | 790    | 143       | 10        | 10,683 |
| % of total voxels      | 84      | 1        | 5        | 7      | 1         | <1        |        |
| <i>White matter</i>    |         |          |          |        |           |           |        |
| Male no. of clusters   | 0       | 1        | 2        | 0      | 0         | 1         | 4      |
| Male total voxels      | 0       | 90       | 517      | 0      | 0         | 25        | 632    |
| % of total voxels      | –       | 14       | 82       | –      | –         | 4         |        |
| Female no. of clusters | 4       | 1        | 4        | 0      | 0         | 2         | 11     |
| Female total voxels    | 4858    | 205      | 527      | 0      | 0         | 31        | 5621   |
| % of total voxels      | 86      | 4        | 9        | –      | –         | <1        |        |

<sup>a</sup> These data are summarized from Tables 1 and 2; sub-clusters are not included.

For GM voxel locations correlated with FSIQ in our study, men exhibited an anterior–posterior combination of bilateral frontal (BA 8, 9) and left parietal (BA 39, 40; Wernicke's area) areas, and women exhibited predominantly frontal lobe regions (BA 10) and, to a lesser extent, Broca's area (BA 44, 45). The role of GM volume in language areas appears to be noteworthy with Broca's area being correlated with FSIQ in women and Wernicke's area in men. Moreover, functional imaging studies with both men and women combined have implicated broad areas of activation during problem solving tasks involving memory and attention, including BA 6, 8, 9, 10, 45, 46, 47 in frontal regions, and BA 7, 39, and 40 in parietal regions (Cabeza and Nyberg, 2000). Thus, our data are consistent with attempts to relate individual differences in general intelligence to fundamental cognitive functions, and support the distributed nature of the neural basis of intelligence throughout the brain (Deary et al., 2004; Gray et al., 2003; Haier et al., 1992, 1988, 2003b; Isaacs et al., 2004) in contrast to the view that the frontal lobes alone are the basis for general intelligence (Duncan et al., 2000).

Women showed fewer GM and more WM voxels related to FSIQ than did men. These results should be interpreted in the context of the sex difference in GM to WM ratio; hence, for each sex, variation in intellectual ability is limited more by the size of voxel clusters in that tissue compartment which is relatively diminished relative to the general human brain design. It is also important to note that functional variation in women's brains emerge from a relatively smaller and more anterior set of clusters. Considered in the context of reduced overall GM and WM volumes in women, and the fact that the sexes do not differ in FSIQ, the current results suggest that different types of brain designs may manifest equivalent intellectual performance.

Other researchers have found that WM sex differences appear to support greater associations with cognition in women than in men. For example, Gur et al. (1999) regressed intelligence on total brain WM and observed that the slope was steeper in women, suggesting that individual variation in WM volume was relatively more important for women than men; this was not the case for total GM. Larger anterior callosal size was significantly correlated with all measures of cognition in women, while no significant correlations were found between regional callosal size and cognitive performance in men

(Davatzikos and Resnick, 1998). Finally, research utilizing Proton Magnetic Resonance Spectroscopy (<sup>1</sup>H-MRS) has established that concentration of N-acetyl-aspartate (NAA) was significantly correlated with a measure of vocabulary (high *g*-loading) in women, but not in men within left frontal WM (Pfleiderer et al., 2004). NAA has been shown in several studies to correlate positively with cognitive ability (Driscoll et al., 2003; Jung et al., 1999a,b; Valenzuela et al., 2000; Yeo et al., 2000). Taken together, these results are consistent with our data indicating relatively more covariation of WM volume and general ability in women.

Interestingly, while early researchers found relatively larger frontal cortices in humans compared to other primates (Brodman, 1912), more contemporary research finds little evidence of proportional differences between primates when great apes are included in the sample (Semendeferi et al., 2002), except for BA 10 which is relatively larger in humans (Semendeferi et al., 2001). Consistent with the current results, both human and animal lesion studies have found that frontal lesions are relatively more detrimental to cognitive performance in women than men. Kimura (1993) noted that for both Block Design and Vocabulary, two WAIS subtests correlating highly with FSIQ, anterior and posterior cerebral lesions had adverse effects for men, but only anterior lesions had a significant adverse effect for women. Similar results emerged with respect to analysis of sex differences in the representation of left hemisphere language skills. Men were rendered aphasic by either anterior or posterior lesions, while anterior lesions were much more apt to produce aphasia than posterior lesions in women. The clusters identified through VBM for each sex in the current analysis thus map onto the left hemisphere regions linked with language for each sex, highlighting the overlap of linguistic and intellectual functions in the human brain.

VBM and ROI methods have their respective strengths and weaknesses (Ashburner and Friston, 2001; Bookstein, 2001; Davatzikos, 2004; Friston and Ashburner, 2004). Some extra caution is required when VBM analysis is applied to relatively small sample sizes like the ones in this study. For this reason, we have limited our discussion mostly to the brain areas identified at  $P < 0.05$ , corrected for multiple comparisons, although this by no means ensures that there are no false-positive results or that all

relevant brain areas have been identified. Also, the mean FSIQ in our samples was about one standard deviation above the population mean. Whether our findings also hold across the full range of FSIQ and a normal distribution must be determined with larger and more representative samples scanned in a standard protocol on the same machine.

The current results contribute to a growing body of evidence demonstrating that, although the sexes do not differ in general intellectual ability, the neural substrates of general intelligence are different. Whether similar neuroanatomical differences are associated with specific mental abilities (assessed for example by WAIS subtests) remains to be determined; our VIQ and PIQ findings need replication with larger samples. Although GM and WM volumes are highly heritable in many areas (Posthuma et al., 2002; Thompson et al., 2001), there is evidence that GM volume in humans can increase with motor learning (Draganski et al., 2004) or the acquisition of a second language (Mechelli et al., 2004) and the nature of genetic determinism in general is now controversial (Silverman, 2004). Future research will require regionally specific analyses of tissue quality (e.g., neurometabolites assessed with MRS, WM matter organization assessed with DTI), coupled with the sorts of morphometric analyses employed here, as well as functional studies of brain activation during a variety of *g* loaded tasks in large samples. Investigation of sex differences may provide a window into the structural and functional brain design principles underlying human intelligence.

#### Acknowledgment

The UCI portion of this work was funded by a grant from NICHD to Dr. Haier (HD037427).

#### References

- Allen, J.S., Damasio, H., Grabowski, T.J., et al., 2003. Sexual dimorphism and asymmetries in the gray-white composition of the human cerebrum. *NeuroImage* 18 (4), 880–894.
- Andreasen, N.C., Flaum, M., Swayze II, V., et al., 1993. Intelligence and brain structure in normal individuals. *Am. J. Psychiatry* 150 (1), 130–144.
- Ashburner, J., Friston, K.J., 2000. Voxel-based morphometry—The methods. *NeuroImage* 11 (6 Pt. 1), 805–821.
- Ashburner, J., Friston, K.J., 2001. Why voxel-based morphometry should be used. *NeuroImage* 14 (6), 1238–1243.
- Bookstein, F.L., 2001. “Voxel-based morphometry” should not be used with imperfectly registered images. *NeuroImage* 14 (6), 1454–1462.
- Brodman, K., 1912. Neue Ergebnisse über die vergleichende histologische Lokalisation der Grosshirnrinde mit besonderer Berücksichtigung des Stirnhirns. *Anat. Anz. (Suppl.)* 41, 157–216.
- Cabeza, R., Nyberg, L., 2000. Imaging cognition II: an empirical review of 275 PET and fMRI studies. *J. Cogn. Neurosci.* 12 (1), 1–47.
- Cahill, L., Haier, R.J., White, N.S., et al., 2001. Sex-related difference in amygdala activity during emotionally influenced memory storage. *Neurobiol. Learn. Mem.* 75 (1), 1–9.
- Davatzikos, C., 2004. Why voxel-based morphometric analysis should be used with great caution when characterizing group differences. *NeuroImage* 23 (1), 17–20.
- Davatzikos, C., Resnick, S.M., 1998. Sex differences in anatomic measures of interhemispheric connectivity: correlations with cognition in women but not men. *Cereb. Cortex* 8 (7), 635–640.
- de Courten-Myers, G.M., 1999. The human cerebral cortex: gender differences in structure and function. *J. Neuropathol. Exp. Neurol.* 58 (3), 217–226.
- Deary, I.J., Simonotto, E., Meyer, M., et al., 2004. The functional anatomy of inspection time: an event-related fMRI study. *NeuroImage* 22 (4), 1466–1479.
- Draganski, B., Gaser, C., Busch, V., et al., 2004. Neuroplasticity: changes in grey matter induced by training. *Nature* 427 (6972), 311–312.
- Driscoll, I., Hamilton, D.A., Petropoulos, H., et al., 2003. The aging hippocampus: cognitive, biochemical and structural findings. *Cereb. Cortex* 13 (12), 1344–1351.
- Duncan, J., Seitz, R.J., Kolodny, J., et al., 2000. A neural basis for general intelligence. *Science* 289 (5478), 457–460.
- Filipek, P.A., Richelme, C., Kennedy, D.N., et al., 1994. The young adult human brain: an MRI-based morphometric analysis. *Cereb. Cortex* 4 (4), 344–360.
- Flashman, L.A., Andreasen, N.C., Flaum, M., et al., 1997. Intelligence and regional brain volumes in normal controls. *Intelligence* 25 (3), 149–160.
- Friston, K., Ashburner, J., 2004. Generative and recognition models for neuroanatomy. *NeuroImage* 23 (1), 21–24.
- Geschwind, N., 1972. Cerebral dominance and anatomic asymmetry. *N. Engl. J. Med.* 287 (4), 194.
- Goldstein, J.M., Seidman, L.J., Horton, N.J., et al., 2001. Normal sexual dimorphism of the adult human brain assessed by in vivo magnetic resonance imaging. *Cereb. Cortex* 11 (6), 490–497.
- Good, C.D., Johnsrude, I.S., Ashburner, J., et al., 2001. A voxel-based morphometric study of ageing in 465 normal adult human brains. *NeuroImage* 14 (1 Pt. 1), 21–36.
- Gray, J.R., Chabris, C.F., Braver, T.S., 2003. Neural mechanisms of general fluid intelligence. *Nat. Neurosci.* 6 (3), 316–322.
- Gur, R.C., Turetsky, B.I., Matsui, M., et al., 1999. Sex differences in brain gray and white matter in healthy young adults: correlations with cognitive performance. *J. Neurosci.* 19 (10), 4065–4072.
- Gur, R.C., Alsop, D., Glahn, D., et al., 2000. An fMRI study of sex differences in regional activation to a verbal and a spatial task. *Brain Lang.* 74 (2), 157–170.
- Haier, R.J., Benbow, C.P., 1995. Sex differences and lateralization in temporal lobe glucose metabolism during mathematical reasoning. *Dev. Neuropsychol.* 11 (4), 405–414.
- Haier, R.J., Siegel, B.V., Nuechterlein, K.H., et al., 1988. Cortical glucose metabolic-rate correlates of abstract reasoning and attention studied with positron emission tomography. *Intelligence* 12 (2), 199–217.
- Haier, R.J., Siegel, B., Tang, C., et al., 1992. Intelligence and changes in regional cerebral glucose metabolic-rate following learning. *Intelligence* 16 (3–4), 415–426.
- Haier, R.J., Alkire, M.T., White, N.S., et al., 2003a. Temporal cortex hypermetabolism in Down syndrome prior to the onset of dementia. *Neurology* 61 (12), 1673–1679.
- Haier, R.J., White, N.S., Alkire, M.T., 2003b. Individual differences in general intelligence correlate with brain function during nonreasoning tasks. *Intelligence* 31 (5), 429–441.
- Haier, R.J., Jung, R., Yeo, R., et al., 2004. Structural brain variation and general intelligence. *NeuroImage* 23 (1), 425–433.
- Isaacs, E.B., Edmonds, C.J., Chong, W.K., et al., 2004. Brain morphometry and IQ measurements in preterm children. *Brain* 127, 2595–2607.
- Jensen, A.R., 1998. The *g* factor: the science of mental ability. Praeger, Westport.
- Jung, R.E., Brooks, W.M., Yeo, R.A., et al., 1999a. Biochemical markers of intelligence: a proton MR spectroscopy study of normal human brain. *Proc. R. Soc. Lond., Ser. B Biol. Sci.* 266 (1426), 1375–1379.
- Jung, R.E., Yeo, R.A., Chiulli, S.J., et al., 1999b. Biochemical markers of cognition: a proton MR spectroscopy study of normal human brain. *NeuroReport* 10 (16), 3327–3331.
- Kimura, D., 1993. *Neuromotor Mechanisms in Human Communication*. Oxford Univ. Press, New York.
- Luders, E., Narr, K.L., Thompson, P.M., et al., 2004. Gender differences in cortical complexity. *Nat. Neurosci.* 7 (8), 799–800.

- Mansour, C.S., Haier, R.J., Buchsbaum, M.S., 1996. Gender comparisons of cerebral glucose metabolic rate in healthy adults during a cognitive task. *Pers. Individ. Differ.* 20 (2), 183–191.
- Mechelli, A., Crinion, J.T., Noppeney, U., et al., 2004. Neurolinguistics: structural plasticity in the bilingual brain. *Nature* 431 (7010), 757.
- Neubauer, A.C., Fink, A., Schrausser, D.G., 2002. Intelligence and neural efficiency: the influence of task content and sex on the brain–IQ relationship. *Intelligence* 30 (6), 515–536.
- Nopoulos, P., Flaum, M., O’Leary, D., et al., 2000. Sexual dimorphism in the human brain: evaluation of tissue volume, tissue composition and surface anatomy using magnetic resonance imaging. *Psychiatry Res.* 98 (1), 1–13.
- Passe, T.J., Rajagopalan, P., Tupler, L.A., et al., 1997a. Age and sex effects on brain morphology. *Prog. Neuro-Psychopharmacol. Biol. Psychiatry* 21 (8), 1231–1237.
- Passe, T.J., Rajagopalan, P., Tupler, L.A., et al., 1997b. Age and sex effects on brain morphology. *Prog. Neuro-Psychopharmacol. Biol. Psychiatry* 21 (8), 1231–1237.
- Peters, M., Jancke, L., Staiger, J.F., et al., 1998. Unsolved problems in comparing brain sizes in *Homo sapiens*. *Brain Cogn.* 37 (2), 254–285.
- Pfleiderer, B., Ohrmann, P., Suslow, T., et al., 2004. N-acetylaspartate levels of left frontal cortex are associated with verbal intelligence in women but not in men: a proton magnetic resonance spectroscopy study. *Neuroscience* 123 (4), 1053–1058.
- Posthuma, D., De Geus, E.J., Baare, W.F., et al., 2002. The association between brain volume and intelligence is of genetic origin. *Nat. Neurosci.* 5 (2), 83–84.
- Price, C.J., Friston, K.J., 1997. Cognitive conjunction: a new approach to brain activation experiments. *NeuroImage* 5 (4 Pt. 1), 261–270.
- Rabinowicz, T., Dean, D.E., Petetot, J.M., et al., 1999. Gender differences in the human cerebral cortex: more neurons in males; more processes in females. *J. Child Neurol.* 14 (2), 98–107.
- Schlaepfer, T.E., Harris, G.J., Tien, A.Y., et al., 1995. Structural differences in the cerebral cortex of healthy female and male subjects: a magnetic resonance imaging study. *Psychiatry Res.* 61 (3), 129–135.
- Semendeferi, K., Armstrong, E., Schleicher, A., et al., 2001. Prefrontal cortex in humans and apes: a comparative study of area 10. *Am. J. Phys. Anthropol.* 114 (3), 224–241.
- Semendeferi, K., Lu, A., Schenker, N., et al., 2002. Humans and great apes share a large frontal cortex prefrontal cortex in humans and apes: a comparative study of area 10. *Nat. Neurosci.* 5 (3), 272–276.
- Shaywitz, B.A., Shaywitz, S.E., Pugh, K.R., et al., 1995. Sex differences in the functional organization of the brain for language. *Nature* 373 (6515), 607–609.
- Shaywitz, B.A., Shaywitz, S.E., Pugh, K.R., et al., 2001. The functional neural architecture of components of attention in language-processing tasks. *NeuroImage* 13 (4), 601–612.
- Silverman, P.H., 2004. Rethinking genetic determinism. *Scientist* 18 (10), 32–33.
- Spearman, C., 1904. General intelligence objectively determined and measured. *Am. J. Psychol.* 15, 201–293.
- Talairach, J., Tournoux, P., 1988. *Co-planar Stereotaxic Atlas of the Human Brain: A 3-dimensional Proportional System. An Approach to Cerebral Imaging.* Stuttgart; New York: New York, G. Thieme; Thieme Medical Publishers.
- Thompson, P.M., Cannon, T.D., Narr, K.L., et al., 2001. Genetic influences on brain structure. *Nat. Neurosci.* 4 (12), 1253–1258.
- Valenzuela, M.J., Sachdev, P.S., Wen, W., et al., 2000. Dual voxel proton magnetic resonance spectroscopy in the healthy elderly: subcortical-frontal axonal N-acetylaspartate levels are correlated with fluid cognitive abilities independent of structural brain changes. *NeuroImage* 12 (6), 747–756.
- Wechsler, D., 1981. *Wechsler Adult Intelligence Scale-revised.* Psychological Corporation, San Antonio, TX.
- Witelson, S.F., Glezer I.I., Kigar, D.L., 1995. Women have greater density of neurons in posterior temporal cortex. *J. Neurosci.* 15 (5 Pt. 1), 3418–3428.
- Yeo, R.A., Hill, D., Campbell, R., et al., 2000. Developmental instability and working memory ability in children: a magnetic resonance spectroscopy investigation. *Dev. Neuropsychol.* 17 (2), 143–159.